AHA FOUNDATION STUDY FINDINGS ON FEMALE GENITAL MUTILATION IN THE U.S.

KENTUCKY

STATE DATA

Based on 2015-2019 American Community Survey population estimates.

8,300 STUDY POPULATON

Women and girls with ancestral ties to countries where FGM/C is practiced

2,083
Women and girls who were likely LIVING
WITH FGM/C

297
Girls who were likely
AT RISK of FGM/C

STATE LEGISLATION AND POLICY LANDSCAPE

STATUS

Strongest Existing Legislation¹

IMPROVE BY ADDING
Nothing

1 https://bit.ly/3RJRzKy



SUMMARY


FGM/C prevalence was estimated at 28.7% within the study population in Kentucky with over 60% of the impacted population in the state identifying as Somali (40.8%), Ethiopian (10.8%) or Egyptian (10.7%).

It is estimated that **601** women were living with Type 3 FGM/C in Kentucky. While all survivors may require some level of medical and mental health support, those living with Type 3 would likely require additional medical attention.

Most of those impacted by FGM/C in Kentucky live in the greater Cincinnati and Louisville/Jefferson County metropolitan areas.

ETHNIC BREAKDOWN

Ethnic breakdown of girls most likely to be AT RISK of FGM/C in Kentucky



NOTE: Nigerian girls are likely underrepresented in this data since they are cut at a very young age, resulting in most girls being encoded as already living with FGM/C.

STATE PREVALENCE RANKING








AGE DISTRIBUTION

Distribution of girls most likely to be AT RISK of FGM/C in Kentucky



SPATIAL DISTRIBUTION

Counties with the highest STUDY POPULATION | LIVING WITH | AT RISK population

Jefferson Fayette	4,550 1,249	1,353 257	226 12
Boone	596	142	16
Warren	326	61	11
Madison	145	52	3
Daviess	54	38	-
Campbell	133	19	3
Kenton	159	17	13
Franklin	50	15	-
Woodford	26	8	-

Metropolitan Areas with the highest STUDY POPULATION | LIVING WITH | AT RISK population

Cincinnati, OH-KY-IN Louisville/Jefferson County, KY-IN Clarksville, TN-KY Owensboro, KY

6,543	1,487	108
4,809	1,422	236
631	167	3
59	41	-

CALL TO ACTION

Interventions tailored to the specifics of the context.

Prevention and response interventions should focus on the greater Cincinnati and Louisville/ Jefferson County metropolitan areas.

Child Protection should focus on **Somali** and **Sudanese** girls between the ages of 5 and 15; **Egyptian** girls between the ages of 6 and 14; **Ethiopian** girls throughout their childhood and adolescence.

All estimates are subject to both sampling and nonsampling error.

For more granular prevalence data contact info@theahafoundation.org

scan to access the full report

